

10/048,212
Updated Search
LYCOOK 11/2/05

d his

(FILE 'HOME' ENTERED AT 14:15:20 ON 02 NOV 2005)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, CANCERLIT, JAPIO' ENTERED AT
14:15:35 ON 02 NOV 2005

L1 3 S (PROTEASE? BOVINE SERUM)
L2 3 DUPLICATE REMOVE L1 (0 DUPLICATES REMOVED)
L3 4334 S (PROTEASE TREAT?)
L4 56 S L3 AND (BOVINE SERUM)
L5 1 S L4 AND AGGLUTIN?
L6 23 DUPLICATE REMOVE L4 (33 DUPLICATES REMOVED)
L7 23 S L6 NOT L5
L8 0 S (PROTEASE TREAT? BOVINE SERUM)
L9 5 S (PROTEASE TREAT? SERUM)
L10 4 DUPLICATE REMOVE L9 (1 DUPLICATE REMOVED)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, CANCERLIT, JAPIO' ENTERED AT
15:07:45 ON 02 NOV 2005

L11 85510 S (BOVINE SERUM ALBUMIN)
L12 657 S L11 AND AGGLUTINATION?
L13 4 S L12 AND PEPSIN?
L14 4 DUPLICATE REMOVE L13 (0 DUPLICATES REMOVED)

d his

(FILE 'HOME' ENTERED AT 14:15:20 ON 02 NOV 2005)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, CANCERLIT, JAPIO' ENTERED AT
14:15:35 ON 02 NOV 2005

L1 3 S (PROTEASE? BOVINE SERUM)
L2 3 DUPLICATE REMOVE L1 (0 DUPLICATES REMOVED)
L3 4334 S (PROTEASE TREAT?)
L4 56 S L3 AND (BOVINE SERUM)
L5 1 S L4 AND AGGLUTIN?
L6 23 DUPLICATE REMOVE L4 (33 DUPLICATES REMOVED)
L7 23 S L6 NOT L5
L8 0 S (PROTEASE TREAT? BOVINE SERUM)
L9 5 S (PROTEASE TREAT? SERUM)
L10 4 DUPLICATE REMOVE L9 (1 DUPLICATE REMOVED)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, CANCERLIT, JAPIO' ENTERED AT
15:07:45 ON 02 NOV 2005

L11 85510 S (BOVINE SERUM ALBUMIN)
L12 657 S L11 AND AGGLUTINATION?
L13 4 S L12 AND PEPSIN?
L14 4 DUPLICATE REMOVE L13 (0 DUPLICATES REMOVED)


[My List - 0](#) [Help](#)
[Search](#)
[Main Search](#) | [Advanced Keyword Search](#) | [Search History](#)
Search:

> You're searching: **Scientific and Technical Information Center**

Item Information

- ▶ [Subscriptions](#)
 - [Holdings](#)
-

Browse Catalog

- by title:
- [International archiv...](#)

MARC Display

International archives of allergy and applied immunology.

Alt. Titles: Proceedings of the International Association of Asthmology.
Author: International Association of Allergists.
Imprint: Basel ; New York : S. Karger, 1950-1991.
Notes: Description based on: Vol. 54, no. 2, published in 1977; title from cover.
 Issued in three volumes per year, four numbers per volume.
 Some volumes accompanied by unnumbered supplements.
 Text in English, French, German or Spanish; summaries in English and the original language.
 Vols. 1-3 issued as official organ of the International Association of Allergists.
 Some issues include: Transactions of the symposia of the Collegium Internationale Allergologicum, or: Proceedings of the International Association of Asthma and of national allergy societies.

ISSN: 0020-5915
Subjects: Allergy -- Periodicals.
 Immunology -- Periodicals.
Description: 96 v. : ill. ; 25 cm.
Continued by: International archives of allergy and immunology

Subscription Summary

US Patent & Trademark Office

Location: US Patent & Trademark Office
Collection: Biotechnology and Chemical Library
Call No.: QR188 .I56
Copy No.: 1
Status: Not Currently Received
Media serial
Type:
Main run: Vol. 105 No. 1 (Sep 1994)
 Vol. 58 No. 1 (1979)- Vol. 87 No. 4 (Dec 1988)

[Show all](#)
 items

US Patent & Trademark Office

Location: US Patent & Trademark Office

Collection: Biotechnology and Chemical Library Microfilm

Call No.: QR188 .I56 Microfilm

Copy No.: 1

Status: Not Currently Received

Media:

film

Type:

Show all
items

Email: pamela.hoeft@uspto.gov to ask questions or make suggestions.

Horizon Information Portal 3.05

Brought to you by *Scientific and Technical Information Center*

ANSWER 4 OF 4 CAPLUS COPYRIGHT 2005 ACS on STN

AN 1969:458970 CAPLUS

DN 71:58970

ED Entered STN: 12 May 1984

TI Red cell-linked antigen-antiglobulin reaction

AU Hunter, A.; Coombs, Robin R. A.

CS Univ. of Cambridge, Cambridge, UK

SO International Archives of Allergy and Applied Immunology (1969), 36,
354-75

CODEN: IAAAAM; ISSN: 0020-5915

DT Journal

LA English

CC 13 (Immunochemistry)

AB Rabbit antibody (I) to human group O erythrocytes (II) was coupled to proteins (III).

Reaction of the I-III complex with II gave a reagent (IV) for determination of antibody to III. Various IV were examined by agglutination titers with rabbit antibody to III and goat antibody to rabbit globulin. Photooxidn. gave fairly good coupling of I to

bovine serum albumin (BSA),

β -lactoglobulin, (β L), castor bean protein, or

α -lactalbumin, but only poor coupling to human serum albumin (HSA), egg albumin, or proteins from pollen extract. These I-III complexes showed little direct agglutination (DA) when used to prepare IV.

Bis-diazobenzidine (BDB) or 2,4-diisocyanotoluene gave better coupling of I to BSA, β L, HSA, or pollen extract, but there was considerable DA.

Neither photooxidn. nor BDB coupled I to PPD of the tubercle bacillus.

Attempts to couple I to BSA with 1-ethyl-3-[3-

(dimethylamino)propyl]carbodiimide gave IV showing excessive clumping.

Coupling of BSA to the 7S fraction of I gave IV showing higher titers than IV prepared with 19S or whole I, but DA was also greater. Coupling of BSA to papain-digested Fab1 fragments of I reduced DA, but only moderate

coupling occurred, and the IV was readily agglutinated by normal rabbit serum.

Coupling of BSA to pepsin-F(ab)2 fragments of I failed

to reduce DA. Coupling of mercaptoethanol-treated pepsin-F(ab)1

fragments reduced DA, but coupling was only moderately good and this IV was agglutinated by a factor difficult to remove from rabbit antisera to human γ -globulin. Ox red cells and rabbit antibody to them could be

substituted for I and II, but without improvement of IV. DA was reduced by photooxidn. of I after coupling it to III by BDB, but the effective

titer of the IV was also reduced; prior photooxidn. gave I which poorly

coupled by BDB to III. Removal by electrophoresis or Sephadex

fractionation of free I from IV reduced but did not eliminate DA.

ST antigens erythrocyte complexes; erythrocyte antigens complexes; antibodies detn reagent; blood typing reagent

IT Antibodies

RL: BIOL (Biological study)

(antigen reactions, erythrocyte protein complexes in)

IT Proteins

RL: BIOL (Biological study)

(erythrocyte complexes, in antibody antigen reactions)

IT Erythrocytes

(protein complexes, in antibody antigen reactions)

10/04 8, 212
updated Search
Lycock 11/2/05

d his

(FILE 'HOME' ENTERED AT 14:15:20 ON 02 NOV 2005)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, CANCERLIT, JAPIO' ENTERED AT
14:15:35 ON 02 NOV 2005

L1 3 S (PROTEASE? BOVINE SERUM)
L2 3 DUPLICATE REMOVE L1 (0 DUPLICATES REMOVED)
L3 4334 S (PROTEASE TREAT?)
L4 56 S L3 AND (BOVINE SERUM)
L5 1 S L4 AND AGGLUTIN?
L6 23 DUPLICATE REMOVE L4 (33 DUPLICATES REMOVED)
L7 23 S L6 NOT L5
L8 0 S (PROTEASE TREAT? BOVINE SERUM)
L9 5 S (PROTEASE TREAT? SERUM)
L10 4 DUPLICATE REMOVE L9 (1 DUPLICATE REMOVED)

=>

d his

(FILE 'HOME' ENTERED AT 14:15:20 ON 02 NOV 2005)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, CANCERLIT, JAPIO' ENTERED AT
14:15:35 ON 02 NOV 2005

L1 3 S (PROTEASE? BOVINE SERUM)
L2 3 DUPLICATE REMOVE L1 (0 DUPLICATES REMOVED)
L3 4334 S (PROTEASE TREAT?)
L4 56 S L3 AND (BOVINE SERUM)
L5 1 S L4 AND AGGLUTIN?
L6 23 DUPLICATE REMOVE L4 (33 DUPLICATES REMOVED)
L7 23 S L6 NOT L5
L8 0 S (PROTEASE TREAT? BOVINE SERUM)
L9 5 S (PROTEASE TREAT? SERUM)
L10 4 DUPLICATE REMOVE L9 (1 DUPLICATE REMOVED)

=>

ANSWER 2 OF 4 CAPLUS COPYRIGHT 2005 ACS on STN

AN 1992:231184 CAPLUS

DN 116:231184

ED Entered STN: 13 Jun 1992

TI Formulation and preliminary testing of a cryptococcal antibody coated latex reagent used with protease pre-treatment

AU Shinoda, Takako; Ikeda, Reiko; Nishikawa, Akemi; Ohtsuka, Morio; Sadamoto, Shinya; Sasaki, Yasuharu; Futami, Shuhei

CS Dep. Microbiol., Meiji Coll. Pharm., Tanashi, 188, Japan

SO Nippon Ishinkin Gakkai Zasshi (1991), 32(Suppl. 2, Proc. Annu. Meet. Jpn. Soc. Med. Mycol., 34th, 1990), 83-93

CODEN: NIGZE4; ISSN: 0916-4804

DT Journal

LA English

CC 9-10 (Biochemical Methods)

Section cross-reference(s): 10, 14

AB The authors defined optimal conditions for detection of cryptococcal antigen using latex particles sensitized with anti-C. neoformans globulin fractionated by 40% saturated ammonium sulfate. Latex particles, with a diameter

of 0.81 μm , were sensitized with 20 μg of anti-C. neoformans globulin per mg latex. The agglutination test was performed using a mixture of 75 μL of **protease treated serum** or

cerebrospinal fluid (CSF) and 25 μL of sensitized latex suspension.

After 10 min reaction on a rotator, the agglutination was read. The authors compared the minimal concentration of polysaccharide antigen detectable with our materials and procedure and with com. available kits and obtained almost the same sensitivities. However, their procedure was also capable of detecting antigen in soluble immune complexes in patient's serum. The sensitivity of their latex agglutination test using the sensitized latex particles was found to be 100% in cases of cryptococcal meningitis, 81.8% in pulmonary cryptococcosis and 75% in cutaneous cryptococcosis. The specificity of this test was 100% with sera and 95% with CSF. Com. kit B was the more useful because of its protease pre-treatment which reduced the problems of false positives due to rheumatoid factor and false neg. due to soluble immune complexes.

ST Cryptococcus antigen latex agglutination test

IT Antigens

RL: ANST (Analytical study)

(cryptococcal, detection of)

IT Cryptococcus neoformans

(detection of, by immunoassay)

IT Immunoassay

(latex agglutination test, for cryptococcal antigens)

IT 9001-92-7, Protease

RL: ANST (Analytical study)

(in cryptococcal antigens detection)